

Publications using Vala's anti-phospho-perilipin 1-serine 522 antibody

Lizardo K, Ayyappan JP, Oswal N, et al. 2021. Fat tissue regulates the pathogenesis and severity of cardiomyopathy in murine chagas disease. *PLoS Negl Trop Dis*; **15**(4):e0008964. doi: 10.1371/journal.pntd.0008964

Depommier C, Van Hul M, Everard A, et al. 2020. Pasteurized *Akkermansia muciniphila* increases whole-body energy expenditure and fecal energy excretion in diet-induced obese mice. *Gut Microbes*; **11**(5):1231-1245. doi: 10.1080/19490976.2020.1737307

Overby H, Yang Y, Xu X, et al. 2020. Soluble epoxide hydrolase inhibition by *t*-TUCB promotes brown adipogenesis and reduces serum triglycerides in diet-induced obesity. *Int J Mol Sci*; **21**(19):E7039. doi: 10.3390/ijms21197039

Krycer JR, Quek L-E, Francis D, et al. 2020. Insulin signaling requires glucose to promote lipid anabolism in adipocytes. *J Biol Chem*; **295**(38):13250-13266. doi: 10.1074/jbc.RA120.014907

Tarabra E, Nouws J, Vash-Margita A, et al. 2020. The omentum of obese girls harbors small adipocytes and browning transcripts. *JCI Insight*; **5**(6):e135448. doi: 10.1172/jci.insight.135448

Ayyappan JP, Lizardo K, Wang S, et al. 2020. Inhibition of SREBP improves cardiac lipidopathy, improves endoplasmic reticulum stress, and modulates chronic Chagas cardiomyopathy. *J Am Heart Assoc*; **9**(3):e014255. doi: 10.1161/JAHA.119.014255

Nayak G, Zhang KX, Vemaraju S, et al. 2020. Adaptive thermogenesis in mice is enhanced by opsin 3-dependent adipocyte light sensing. *Cell Rep*; **30**(3):672-686. doi: 10.1016/j.celrep.2019.12.043

Yang Y, Fu M, Li M-D, et al. 2020. O-GlcNAc transferase inhibits visceral fat lipolysis and promotes diet-induced obesity. *Nat Commun*; **11**(1):181. doi: 10.1038/s41467-019-13914-8

El Ouarrat D, Isaac R, Lee YS, et al. 2020. TAZ is a negative regulator of PPAR γ activity in adipocytes and TAZ deletion improves insulin sensitivity and glucose tolerance. *Cell Metab*; **31**(1):162-173. doi: 10.1016/j.cmet.2019.10.003

Lowry JE, Tumurbaatar B, D'Agostino C, et al. 2019. Effect of high-fat diet on peripheral blood mononuclear cells and adipose tissue in early stages of diet-induced weight gain. *Br J Nutr*; **122**(2):1359-1367. doi: 10.1017/S0007114519002472

Jung SM, Hung C-M, Hildebrand SR, et al. 2019. Non-canonical mTORC2 signaling regulates brown adipocyte lipid catabolism through SIRT6-FoxO1. *Mol Cell*; **75**(4):807-822. doi: 10.1016/j.molcel.2019.07.023

Abulizi A, Camporez J-P, Jurczak MJ, et al. 2019. Adipose glucocorticoid action influences whole-body metabolism via modulation of hepatic insulin action. *FASEB J*; **33**(7):8174-8185. doi: 10.1096/fj.201802706R

Krois CR, Vuckovic MG, Huang P, et al. 2019. RDH1 suppresses adiposity by promoting brown adipose adaptation to fasting and re-feeding. *Cell Mol Life Sci*; **76**(12):2425-2447. doi: 10.1007/s00018-019-03046-z

- Liu K, Yu W, Wei W, et al. 2019. Melatonin reduces intramuscular fat deposition by promoting lipolysis and increasing mitochondrial function. *J Lipid Res*; **60**(4):767-782. doi: 10.1194/jlr.M087619
- Boudreau A, Richard AJ, Burrell JA, et al. 2018. An ethanolic extract of *Artemisia scoparia* inhibits lipolysis in vivo and has antilipolytic effects on murine adipocytes in vitro. *Am J Physiol Endocrinol Metab*; **315**(5):E1053-E1061. doi: 10.1152/ajpendo.00177.2018
- Hashani M, Witzel HR, Pawella LM, et al. 2018. Widespread expression of perilipin 5 in normal human tissues and in diseases is restricted to distinct lipid droplet subpopulations. *Cell Tissue Res*; **374**(1):121-136. doi: 10.1007/s00441-018-2845-7
- Rogne M, Chu D-T, Küntzinger TM, et al. 2018. OPA1-anchored PKA phosphorylates perilipin 1 on S522 and S497 in adipocytes differentiated from human adipose stem cells. *Mol Biol Cell*; **29**(12):1487-1501. doi: 10.1091/mbc.E17-09-0538
- Sanchez-Gurmanches J, Tang Y, Jespersen NZ, et al. 2018. Brown fat AKT2 is a cold-induced kinase that stimulates ChREBP-mediated de novo lipogenesis to optimize fuel storage and thermogenesis. *Cell Metab*; **27**(1):195-209. doi: 10.1016/j.cmet.2017.10.008
- Rondini EA, Mladenovic-Lucas L, Roush WR, et al. 2017. Novel pharmacological probes reveal ABHD5 as a locus of lipolysis control in white and brown adipocytes. *J Pharmacol Exp Ther*; **363**(3):367-376. doi: 10.1124/jpet.117.243253
- Hansen JS, de Maré S, Jones HA, et al. 2017. Visualization of lipid directed dynamics of perilipin 1 in human primary adipocytes. *Sci Rep*; **7**(1):15011. doi: 10.1038/s41598-017-15059-4
- Guilherme A, Pedersen DJ, Henchey E, et al. 2017. Adipocyte lipid synthesis coupled to neuronal control of thermogenic programming. *Mol Metab*; **6**(8):781-796. doi: 10.1016/j.molmet.2017.05.012
- Rui Y, Tong L, Cheng J, et al. 2017. Rosmarinic acid suppresses adipogenesis, lipolysis in 3T3-L1 adipocytes, lipopolysaccharide-stimulated tumor necrosis factor- α secretion in macrophages, and inflammatory mediators in 3T3-L1 adipocytes. *Food Nutr Res*; **61**(1):1330096. doi: 10.1080/16546628.2017.1330096
- Kuo T, Chen T-C, Lee RA, et al. 2017. Pik3r1 is required for glucocorticoid-induced perilipin 1 phosphorylation in lipid droplet for adipocyte lipolysis. *Diabetes*; **66**(6):1601-1610. doi: 10.2337/db16-0831
- Tumurbaatar B, Poole AT, Olson G, et al. 2017. Adipose tissue insulin resistance in gestational diabetes. *Metab Syndr Relat Disord*; **15**(2):86-92. doi: 10.1089/met.2016.0124
- Keskin I, Sutcu M, Eren H, Keskin M. 2016. Exposure to tumescent solution significantly increases phosphorylation of perilipin in adipocytes. *Aesthet Surg J*; **37**(2):239-245. doi: 10.1093/asj/sjw156
- Ding L, Zhang F, Zhao M-X, et al. 2016. Reduced lipolysis response to adipose afferent reflex involved in impaired activation of adrenoceptor-cAMP-PKA-hormone sensitive lipase pathway in obesity. *Sci Rep*; **6**():34374. doi: 10.1038/srep34374

- Krishnamoorthy L, Cotruvo JA Jr., Chang H, et al. 2016. Copper regulates cyclic AMP-dependent lipolysis. *Nat Chem Biol*; **12**(8):586-592. doi: 10.1038/nchembio.2098
- Sagar G, Sah RP, SK Javeed, et al. 2015. Pathogenesis of pancreatic cancer exosome-induced lipolysis in adipose tissue. *Gut*; **65**(7):1165-1174. doi: 10.1136/gutjnl-2014-308350
- Larsson S, Jones HA, Göransson O, et al. 2015. Parathyroid hormone induces adipocyte lipolysis via PKA-mediated phosphorylation of hormone-sensitive lipase. *Cell Signal*; **28**(3):204-213. doi: 10.1016/j.cellsig.2015.12.012
- Sanders MA, Madoux F, Mladenovic H, et al. 2015. Endogenous and synthetic ABHD5 ligands regulate ABHD5-perilipin interactions and lipolysis in fat and muscle. *Cell Metab*; **22**(5):851-860. doi: 10.1016/j.cmet.2015.08.023
- Martins GPC, Souza CO, da O Marques S, et al. 2015. Topiramate effects lipolysis in 3T3-L1 adipocytes. *Biomed Rep*; **3**(6):827-830. doi: 10.3892/br.2015.514
- DiPilato LM, Ahmad F, Harms M, et al. 2015. The role of PDE3B phosphorylation in the inhibition of lipolysis by insulin. *Mol Cell Biol*; **35**(16):2752-2760. doi: 10.1128/MCB.00422-15
- Wiedemann MSF, Wueest S, Grob A, et al. 2014. Short-term HFD does not alter lipolytic function of adipocytes. *Adipocyte*; **3**(2):115-120. doi: 10.4161/adip.27575
- Mowers J, Uhm M, Reilly SM, et al. 2013. Inflammation produces catecholamine resistance in obesity via activation of PDE3B by the protein kinases IKK ϵ and TBK1. *eLife*; **2**:e01119. doi: 10.7554/eLife.01119
- McDonough PM, Maciejewski-Lenoir D, Hartig SM, et al. 2013. Differential phosphorylation of perilipin 1A at the initiation of lipolysis revealed by novel monoclonal antibodies and high content analysis. *PLoS ONE*; **8**(2):e55511. doi: 10.1371/journal.pone.0055511
- Rapold RA, Wueest S, Knoepfel A, et al. 2013. Fas activates lipolysis in a Ca²⁺-CaMKII-dependent manner in 3T3-L1 adipocytes. *J Lipid Res*; **54**(1):63-70. doi: 10.1194/jlr.M028035
- Singh V, Jamwal S, Jain R, et al. 2012. Mycobacterium tuberculosis-driven targeted recalibration of macrophage lipid homeostasis promotes the foamy phenotype. *Cell Host & Microbe*; **12**(5):669-681. doi: 10.1016/j.chom.2012.09.012
- Zhou D, Samovski D, Okunade AL, et al. 2012. CD36 level and trafficking are determinants of lipolysis in adipocytes. *FASEB J*; **26**(11):4733-4742. doi: 10.1096/fj.12-206862
- Nguyen CD, Qiu Y, Cui X, et al. 2011. Alternatively activated macrophages produce catecholamines to sustain adaptive thermogenesis. *Nature*; **480**(7375):104-108. doi: 10.1038/nature10653